

2012

# 5a-Butyl-1,3,8,10-tetra-chloro-7,13-bis-(4-nitro-benzo-yl)-5a,6a,12a,12b-tetra-hydro-7H,13H-thieno[2,3-b:4,5-b']bis-(1,4-benzoxazine)

Kai Tang  
*Bucknell University*

Margaret E. Kastner  
*Bucknell University*, [kastner@bucknell.edu](mailto:kastner@bucknell.edu)

Follow this and additional works at: [http://digitalcommons.bucknell.edu/fac\\_journ](http://digitalcommons.bucknell.edu/fac_journ)

 Part of the [Chemistry Commons](#)

---

## Recommended Citation

Tang, Kai and Kastner, Margaret E.. "5a-Butyl-1,3,8,10-tetra-chloro-7,13-bis-(4-nitro-benzo-yl)-5a,6a,12a,12b-tetra-hydro-7H,13H-thieno[2,3-b:4,5-b']bis-(1,4-benzoxazine)." *Acta crystallographica. Section E, Structure reports online* 68, no. Pt 9 (2012) : o2647-o2648.

This Article is brought to you for free and open access by the Faculty Research and Publications at Bucknell Digital Commons. It has been accepted for inclusion in Faculty Journal Articles by an authorized administrator of Bucknell Digital Commons. For more information, please contact [dcadmin@bucknell.edu](mailto:dcadmin@bucknell.edu).

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 5a-Butyl-1,3,8,10-tetrachloro-7,13-bis(4-nitrobenzoyl)-5a,6a,12a,12b-tetrahydro-7H,13H-thieno[2,3-b:4,5-b']bis(1,4-benzoxazine)

Kai Tang and Margaret E. Kastner\*

Department of Chemistry, Bucknell University, Lewisburg, PA 17837, USA

Correspondence e-mail: kastner@bucknell.edu


Received 24 July 2012; accepted 31 July 2012

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å; disorder in main residue;  $R$  factor = 0.045;  $wR$  factor = 0.123; data-to-parameter ratio = 17.3.

The title compound,  $\text{C}_{34}\text{H}_{24}\text{Cl}_4\text{N}_4\text{O}_8\text{S}$ , is a linear pentacyclic system formed of two substituted benzoxazinyl groups fused to 2-*n*-butyltetrahydrothiophene. The oxazine ring, which is fused to the *n*-butyl-substituted side of the thiophene ring, is in a boat conformation. The other fused oxazine ring and the tetrahydrothiophene ring are each in an envelope conformation. The bridgehead C atom  $\alpha$  to both the S and N atoms forms the flap of each envelope. This results in a twist of the pentacyclic system such that the dihedral angle between the terminal dichlorobenzene rings is  $82.92(8)^\circ$ . In the crystal, inversion-related molecules form a weakly hydrogen-bonded dimer, with two  $\text{C}-\text{H}\cdots\text{O}$  interactions between an H atom on the oxazine ring and an amide O atom. Additionally,  $\text{C}-\text{H}\cdots\text{O}$  interactions occur between an H atom on a screw-related nitrobenzene ring and an O atom on the nitrobenzene ring of one molecule. One of the Cl atoms and the butyl group are disordered over two sets of sites with occupancy ratios of 0.94 (2):0.06 (2) and 0.624 (4):0.376 (4), respectively.

## Related literature

For the synthesis of the title compound, see: Heine *et al.* (1993). For the crystal structure of a related compound, see: Garbaskas *et al.* (1985).



## Experimental

### Crystal data

$\text{C}_{34}\text{H}_{24}\text{Cl}_4\text{N}_4\text{O}_8\text{S}$   
 $M_r = 790.44$   
 Monoclinic,  $P2_1/c$   
 $a = 12.855(3)$  Å  
 $b = 10.139(1)$  Å  
 $c = 27.845(5)$  Å  
 $\beta = 100.23(2)^\circ$

$V = 3571.5(11)$  Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.45$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.5 \times 0.2 \times 0.1$  mm

### Data collection

Siemens R3m/V diffractometer  
 8579 measured reflections  
 8215 independent reflections  
 5039 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.032$   
 3 standard reflections every 97 reflections  
 intensity decay: none

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.123$   
 $S = 1.01$   
 8215 reflections  
 476 parameters

6 restraints  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.34$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.32$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}2-\text{H}2\cdots\text{O}2^i$	0.98	2.55	3.491 (3)	162
$\text{C}14-\text{H}14\cdots\text{O}3^{ii}$	0.93	2.41	3.298 (4)	160

Symmetry codes: (i)  $-x + 1, -y + 1, -z + 1$ ; (ii)  $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$ .

Data collection: XSCANS (Siemens, 1996); cell refinement: XSCANS; data reduction: XSCANS; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors thank the National Science Foundation for grant # ILL8951058 for financial assistance.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2572).

## References

- Garbaskas, M. F., Williams, E. A. & Heine, H. W. (1985). *Acta Cryst.* **C41**, 1217–1222.
- Heine, H. W., Williams, D. K., Rutherford, J. L., Ramphal, J. & Williams, E. A. (1993). *Heterocycles*, **35**, 1125–1140.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Siemens (1996). *XSCANS*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

## supplementary materials

*Acta Cryst.* (2012). E68, o2647–o2648 [doi:10.1107/S1600536812034149]

**5a-Butyl-1,3,8,10-tetrachloro-7,13-bis(4-nitrobenzoyl)-5a,6a,12a,12b-tetrahydro-7H,13H-thieno[2,3-b:4,5-b']bis(1,4-benzoxazine)****Kai Tang and Margaret E. Kastner****Comment**

The title compound was synthesized as part of a study of inverse electron-demand Diels-Alder reactions (Heine *et al.*, 1993).

The title molecule (Fig. 1), it is a linear pentacyclic system formed of two substituted benzoxazinyl groups fused to 2-n-butyltetrahydrothiophene. The oxazine ring (N1/O1/C1–C4) is in a boat conformation with O1 and N1 displaced by 0.536 (3) and 0.478 (3) Å, respectively, from the plane of C1–C4 atoms (r.m.s.d = 0.019 Å). In the dichlorobenzoxazinyl ring (O1/N2/C11–C8/C11/C12), the atoms O1/N1/C3–C8 lie in a plane (rmsd 0.0278 Å) while the atoms C1, C2, C11 and C12 lie 1.035 (3), 0.960 (3), 0.269 (2) and 0.014 (2) Å, respectively, out of this plane. In the other dichlorobenzoxazinyl ring (O5/N3/C16–C23/C13/C14), the atoms N3, O5, C16 and C18–C23, are coplanar (r.m.s.d = 0.0289 Å) while the atoms C17, C13, C14A and C14B are displaced from this plane by 0.553 (3), 0.170 (2), 0.52 (7) and 0.003 (8) Å, respectively. The dihedral angle between the terminal dichlorobenzene rings is 82.92 (8)°. The tetrahydrothiophene ring (S1/C1/C2/C16/C17) is in a C17-envelope conformation with C17 displaced by 0.630 (3) Å from the plane of the remaining ring atoms (r.m.s.d = 0.042 Å). The oxazine ring (N3/O5/C16–C19) is also in a C17-envelope conformation with C17 displaced by 0.572 (3) Å from the plane the other five atoms in the ring (r.m.s.d = 0.035 Å). The dihedral angles between the nitro groups and their associated benzene rings are: 5.9 (5)° for N/O3/O4 and 14.1 (3)° for N4/O7/O8.

In the crystal two inversion-related molecules form a weakly hydrogen-bonded dimer with C2—H2...O2 interactions (Table 1 and Fig. 2) Additionally, C14—H14...O3 hydrogen bonding interactions form between an oxygen on the nitrobenzene ring of one molecule and a hydrogen on a screw related nitrobenzene ring.

**Experimental**

The title compound was synthesized by following a reported procedure (Heine *et al.*, (1993)). The crystals suitable for X-ray crystallographic analysis were grown from a solution of acetonitrile by slow evaporation at room temperature.


**Refinement**

The H atoms were positioned geometrically and refined using a riding model, with C—H = 0.93, 0.96, 0.97 and 0.98 Å, for aryl, methyl, methylene and methyne H-atoms, respectively. The  $U_{iso}(H)$  were allowed at  $1.5U_{eq}(C \text{ methyl})$  or  $1.2U_{eq}(C \text{ non-methyl})$ .

The *n*-butyl group was disordered [ratio: 0.624 (4):0.376 (4)] and the bond distances were constrained to chemically acceptable values. A chlorine atom, Cl4, was also disordered with a population ratio of 0.94 (2):0.06 (2).


### Computing details

Data collection: *XSCANS* (Siemens, 1996); cell refinement: *XSCANS* (Siemens, 1996); data reduction: *XSCANS* (Siemens, 1996); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).




**Figure 1**

The molecular structure of the title compound, with atom labels and 30% probability displacement ellipsoids for non-H atoms. Cl4 and the *n*-butyl group are disordered and only one position is shown.




**Figure 2**

A view of the C—H...O hydrogen bonds (dotted lines) in the crystal structure of the title compound. H atoms non-participating in hydrogen-bonding were omitted for clarity.


**Figure 3**

The molecular structure of the title compound with labels for the disordered atoms and 30% probability displacement ellipsoids for non-H atoms.



**Figure 4**

An enhanced figure of the molecule.


**Figure 5**

An enhanced figure of the unit cell.

**5a-Butyl-1,3,8,10-tetrachloro-7,13-bis(4-nitrobenzoyl)-5a,6a,12a,12b-tetrahydro-7H,13H-thieno[2,3-b:4,5-b']bis(1,4-benzoxazine)**

*Crystal data*

$C_{34}H_{24}Cl_4N_4O_8S$

$M_r = 790.44$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 12.855 (3) \text{ \AA}$

$b = 10.139 (1) \text{ \AA}$

$c = 27.845 (5) \text{ \AA}$

$\beta = 100.23 (2)^\circ$

$V = 3571.5 (11) \text{ \AA}^3$

$Z = 4$

$F(000) = 1616$

$D_x = 1.470 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 20 reflections

$\theta = 10\text{--}12.5^\circ$

$\mu = 0.45 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Needle, colorless

$0.5 \times 0.2 \times 0.1 \text{ mm}$

*Data collection*

Siemens R3m/V

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\theta$  2  $\theta$  scans

8579 measured reflections

8215 independent reflections

5039 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.032$

$\theta_{\text{max}} = 27.6^\circ$ ,  $\theta_{\text{min}} = 1.6^\circ$

$h = 0 \rightarrow 16$

$k = 0 \rightarrow 13$

$l = -36 \rightarrow 35$

3 standard reflections every 97 reflections

intensity decay: none

*Refinement*

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.045$

$wR(F^2) = 0.123$

$S = 1.00$

8215 reflections

476 parameters

6 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0624P)^2]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} = 0.017$

$\Delta\rho_{\text{max}} = 0.34 \text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.32 \text{ e \AA}^{-3}$



*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C11	0.86026 (6)	0.33223 (7)	0.62869 (3)	0.05638 (19)	
C12	0.95304 (6)	0.70359 (9)	0.76649 (3)	0.0707 (2)	
C13	1.04827 (5)	0.89267 (7)	0.45592 (2)	0.05211 (18)	
C14A	0.704 (3)	0.903 (6)	0.3231 (17)	0.1013 (10)	0.06 (2)
C14B	0.7177 (2)	0.8654 (5)	0.31026 (17)	0.1013 (10)	0.94 (2)
S	0.76688 (6)	0.84957 (6)	0.55950 (2)	0.04857 (18)	
O1	0.64969 (13)	0.75499 (16)	0.62293 (5)	0.0394 (4)	
O2	0.57552 (17)	0.37288 (18)	0.53786 (6)	0.0611 (5)	
O3	0.5516 (3)	−0.0704 (3)	0.71927 (11)	0.1308 (13)	
O4	0.5328 (3)	0.0893 (3)	0.76668 (10)	0.1139 (11)	
O5	0.70106 (13)	0.65736 (18)	0.47193 (5)	0.0461 (4)	
O6	1.06030 (14)	0.62308 (17)	0.49661 (6)	0.0481 (4)	
O7	1.2038 (3)	0.5453 (3)	0.74913 (9)	0.1113 (10)	
O8	1.2033 (2)	0.7573 (3)	0.75138 (8)	0.0896 (8)	
N1	0.67147 (14)	0.51822 (18)	0.58817 (6)	0.0334 (4)	
N2	0.5500 (2)	0.0458 (3)	0.72858 (11)	0.0791 (8)	
N3	0.91180 (15)	0.72470 (19)	0.51363 (6)	0.0373 (4)	
N4	1.1870 (2)	0.6530 (3)	0.72987 (10)	0.0715 (7)	
C1	0.65006 (19)	0.7586 (2)	0.57105 (8)	0.0369 (5)	
C2	0.65631 (18)	0.6171 (2)	0.54903 (7)	0.0325 (5)	
H2	0.5894	0.5986	0.5271	0.039*	
C3	0.72924 (18)	0.6790 (2)	0.64898 (8)	0.0357 (5)	
C4	0.74216 (17)	0.5532 (2)	0.63131 (8)	0.0335 (5)	
C5	0.82759 (19)	0.4781 (2)	0.65429 (8)	0.0392 (5)	
C6	0.8903 (2)	0.5241 (3)	0.69675 (9)	0.0460 (6)	
H6	0.9452	0.4726	0.7132	0.055*	
C7	0.8704 (2)	0.6469 (3)	0.71427 (8)	0.0471 (6)	
C8	0.7918 (2)	0.7280 (3)	0.69062 (8)	0.0441 (6)	
H8	0.7813	0.8121	0.7022	0.053*	
C9	0.6174 (2)	0.4013 (2)	0.57913 (8)	0.0394 (5)	
C10	0.6058 (2)	0.3136 (2)	0.62103 (9)	0.0403 (6)	
C11	0.6156 (2)	0.1783 (2)	0.61429 (10)	0.0538 (7)	
H11	0.6343	0.1471	0.5856	0.065*	
C12	0.5980 (2)	0.0908 (3)	0.64951 (11)	0.0592 (8)	
H12	0.6059	0.0005	0.6455	0.071*	
C13	0.5685 (2)	0.1405 (3)	0.69056 (10)	0.0509 (7)	
C14	0.5575 (2)	0.2728 (3)	0.69851 (10)	0.0533 (7)	

H14	0.5375	0.3030	0.7271	0.064*	
C15	0.5767 (2)	0.3602 (2)	0.66308 (9)	0.0453 (6)	
H15	0.5699	0.4505	0.6676	0.054*	
C16	0.74536 (17)	0.6109 (2)	0.51966 (7)	0.0335 (5)	
H16	0.7711	0.5202	0.5182	0.040*	
C17	0.83346 (18)	0.7009 (2)	0.54434 (8)	0.0351 (5)	
H17	0.8684	0.6591	0.5747	0.042*	
C18	0.76444 (19)	0.7222 (2)	0.44547 (8)	0.0403 (6)	
C19	0.86932 (19)	0.7570 (2)	0.46436 (8)	0.0368 (5)	
C20	0.92359 (19)	0.8300 (2)	0.43419 (9)	0.0410 (6)	
C21	0.8790 (2)	0.8600 (3)	0.38661 (9)	0.0516 (7)	
H21	0.9178	0.9044	0.3665	0.062*	
C22	0.7756 (2)	0.8230 (3)	0.36943 (9)	0.0587 (8)	
C23	0.7176 (2)	0.7562 (3)	0.39837 (9)	0.0538 (7)	
H23	0.6476	0.7340	0.3865	0.065*	
C24	1.01125 (19)	0.6649 (2)	0.52636 (9)	0.0392 (5)	
C25	1.05379 (19)	0.6615 (2)	0.58040 (9)	0.0422 (6)	
C26	1.0972 (2)	0.5471 (3)	0.60153 (10)	0.0582 (8)	
H26	1.0970	0.4715	0.5826	0.070*	
C27	1.1411 (2)	0.5429 (3)	0.65061 (11)	0.0658 (8)	
H27	1.1695	0.4651	0.6650	0.079*	
C28	1.1417 (2)	0.6559 (3)	0.67727 (9)	0.0535 (7)	
C29	1.0998 (2)	0.7720 (3)	0.65746 (10)	0.0618 (8)	
H29	1.1024	0.8479	0.6764	0.074*	
C30	1.0537 (2)	0.7742 (3)	0.60879 (10)	0.0584 (8)	
H30	1.0225	0.8512	0.5950	0.070*	
C31A	0.5497 (2)	0.8317 (3)	0.54905 (9)	0.0552 (7)	
H31A	0.5615	0.9252	0.5549	0.066*	0.624 (4)
H31B	0.5374	0.8183	0.5140	0.066*	0.624 (4)
C32A	0.4516 (5)	0.7932 (9)	0.5675 (4)	0.074 (2)	0.624 (4)
H32A	0.4621	0.8011	0.6028	0.089*	0.624 (4)
H32B	0.4320	0.7030	0.5585	0.089*	0.624 (4)
C33A	0.3653 (4)	0.8902 (6)	0.5431 (2)	0.0687 (13)	0.624 (4)
H33A	0.3861	0.9804	0.5515	0.082*	0.624 (4)
H33B	0.3544	0.8810	0.5079	0.082*	0.624 (4)
C34A	0.2662 (4)	0.8563 (8)	0.5617 (3)	0.104 (2)	0.624 (4)
H34A	0.2399	0.7728	0.5485	0.156*	0.624 (4)
H34B	0.2139	0.9232	0.5518	0.156*	0.624 (4)
H34C	0.2811	0.8511	0.5966	0.156*	0.624 (4)
H31C	0.5451	0.8378	0.5140	0.066*	0.376 (4)
H31D	0.5522	0.9205	0.5621	0.066*	0.376 (4)
C32B	0.4516 (7)	0.7606 (17)	0.5604 (7)	0.074 (2)	0.376 (4)
H32C	0.4613	0.7552	0.5957	0.089*	0.376 (4)
H32D	0.4561	0.6709	0.5489	0.089*	0.376 (4)
C33B	0.3355 (7)	0.8037 (9)	0.5432 (4)	0.0687 (13)	0.376 (4)
H33C	0.3209	0.8099	0.5079	0.082*	0.376 (4)
H33D	0.2885	0.7381	0.5530	0.082*	0.376 (4)
C34B	0.3156 (9)	0.9355 (11)	0.5651 (5)	0.104 (2)	0.376 (4)
H34D	0.3310	0.9295	0.6000	0.156*	0.376 (4)

H34E	0.2429	0.9598	0.5548	0.156*	0.376 (4)
H34F	0.3603	1.0010	0.5543	0.156*	0.376 (4)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl1	0.0574 (4)	0.0453 (4)	0.0640 (4)	0.0154 (3)	0.0044 (3)	-0.0023 (3)
Cl2	0.0673 (5)	0.0899 (6)	0.0467 (4)	-0.0176 (4)	-0.0120 (3)	-0.0106 (4)
Cl3	0.0555 (4)	0.0480 (4)	0.0555 (4)	-0.0114 (3)	0.0173 (3)	0.0002 (3)
Cl4A	0.0683 (7)	0.195 (2)	0.0404 (11)	0.0100 (9)	0.0105 (7)	0.0446 (13)
Cl4B	0.0683 (7)	0.195 (2)	0.0404 (11)	0.0100 (9)	0.0105 (7)	0.0446 (13)
S	0.0659 (5)	0.0321 (3)	0.0518 (4)	-0.0050 (3)	0.0216 (3)	-0.0053 (3)
O1	0.0462 (9)	0.0421 (9)	0.0305 (8)	0.0102 (8)	0.0081 (7)	-0.0036 (7)
O2	0.0892 (15)	0.0464 (11)	0.0408 (10)	-0.0161 (10)	-0.0071 (10)	-0.0045 (8)
O3	0.229 (4)	0.0590 (17)	0.125 (2)	-0.021 (2)	0.085 (3)	0.0261 (16)
O4	0.194 (3)	0.092 (2)	0.0727 (16)	0.003 (2)	0.0686 (19)	0.0203 (15)
O5	0.0429 (10)	0.0654 (12)	0.0296 (8)	-0.0093 (9)	0.0054 (7)	0.0029 (8)
O6	0.0498 (11)	0.0483 (10)	0.0480 (10)	0.0046 (9)	0.0132 (8)	-0.0026 (8)
O7	0.147 (3)	0.102 (2)	0.0679 (16)	0.0114 (19)	-0.0277 (16)	0.0222 (15)
O8	0.0913 (18)	0.106 (2)	0.0600 (14)	0.0006 (15)	-0.0163 (12)	-0.0223 (14)
N1	0.0385 (11)	0.0317 (10)	0.0291 (9)	-0.0023 (8)	0.0038 (8)	-0.0002 (8)
N2	0.102 (2)	0.069 (2)	0.0735 (19)	-0.0069 (17)	0.0343 (17)	0.0208 (16)
N3	0.0373 (11)	0.0438 (11)	0.0303 (10)	-0.0007 (9)	0.0047 (8)	0.0020 (8)
N4	0.0691 (18)	0.087 (2)	0.0508 (15)	0.0045 (16)	-0.0092 (13)	0.0002 (15)
C1	0.0467 (14)	0.0328 (12)	0.0320 (11)	0.0063 (11)	0.0094 (10)	-0.0013 (9)
C2	0.0356 (12)	0.0309 (11)	0.0302 (11)	0.0005 (10)	0.0041 (9)	-0.0007 (9)
C3	0.0401 (13)	0.0385 (13)	0.0291 (11)	-0.0004 (10)	0.0082 (9)	-0.0006 (9)
C4	0.0379 (13)	0.0339 (12)	0.0288 (11)	-0.0028 (10)	0.0059 (9)	0.0007 (9)
C5	0.0409 (14)	0.0358 (13)	0.0403 (13)	0.0009 (11)	0.0061 (10)	0.0049 (10)
C6	0.0436 (15)	0.0502 (15)	0.0410 (14)	-0.0003 (12)	-0.0018 (11)	0.0096 (12)
C7	0.0487 (15)	0.0562 (16)	0.0334 (12)	-0.0095 (13)	-0.0006 (11)	-0.0007 (11)
C8	0.0529 (15)	0.0437 (14)	0.0354 (13)	-0.0046 (12)	0.0068 (11)	-0.0051 (11)
C9	0.0471 (14)	0.0333 (12)	0.0376 (13)	-0.0004 (11)	0.0067 (11)	-0.0038 (10)
C10	0.0456 (14)	0.0353 (13)	0.0399 (13)	-0.0074 (11)	0.0078 (11)	-0.0019 (10)
C11	0.083 (2)	0.0342 (14)	0.0486 (15)	-0.0082 (13)	0.0236 (14)	-0.0053 (11)
C12	0.085 (2)	0.0327 (14)	0.0626 (18)	-0.0065 (14)	0.0212 (16)	0.0009 (13)
C13	0.0582 (17)	0.0443 (15)	0.0527 (16)	-0.0083 (13)	0.0166 (13)	0.0093 (12)
C14	0.0617 (18)	0.0567 (17)	0.0469 (15)	-0.0014 (14)	0.0245 (13)	-0.0017 (13)
C15	0.0530 (16)	0.0354 (13)	0.0503 (15)	-0.0003 (12)	0.0166 (12)	-0.0025 (11)
C16	0.0405 (13)	0.0317 (11)	0.0294 (11)	0.0003 (10)	0.0090 (9)	-0.0007 (9)
C17	0.0409 (13)	0.0343 (12)	0.0305 (11)	-0.0021 (10)	0.0072 (10)	0.0004 (9)
C18	0.0436 (14)	0.0474 (14)	0.0318 (12)	0.0003 (11)	0.0120 (10)	-0.0017 (10)
C19	0.0425 (13)	0.0384 (13)	0.0302 (11)	0.0037 (11)	0.0084 (10)	-0.0001 (10)
C20	0.0459 (14)	0.0395 (13)	0.0409 (13)	0.0047 (11)	0.0168 (11)	0.0014 (11)
C21	0.0589 (18)	0.0593 (17)	0.0406 (14)	0.0050 (14)	0.0196 (12)	0.0096 (12)
C22	0.0577 (18)	0.087 (2)	0.0319 (13)	0.0132 (16)	0.0105 (12)	0.0116 (14)
C23	0.0450 (15)	0.081 (2)	0.0350 (13)	0.0040 (14)	0.0062 (11)	0.0062 (13)
C24	0.0400 (13)	0.0312 (12)	0.0454 (13)	-0.0050 (11)	0.0042 (11)	0.0010 (10)
C25	0.0373 (13)	0.0427 (14)	0.0447 (13)	-0.0008 (11)	0.0022 (11)	0.0000 (11)

C26	0.0675 (19)	0.0477 (16)	0.0551 (17)	0.0168 (14)	-0.0011 (14)	-0.0082 (13)
C27	0.073 (2)	0.0572 (18)	0.0605 (18)	0.0225 (16)	-0.0067 (16)	0.0054 (15)
C28	0.0487 (16)	0.0618 (18)	0.0453 (15)	0.0036 (14)	-0.0040 (12)	0.0029 (13)
C29	0.077 (2)	0.0492 (17)	0.0542 (17)	-0.0015 (15)	-0.0024 (15)	-0.0121 (14)
C30	0.074 (2)	0.0413 (15)	0.0522 (16)	0.0025 (14)	-0.0095 (14)	-0.0007 (13)
C31A	0.0674 (19)	0.0580 (17)	0.0371 (13)	0.0326 (15)	0.0008 (12)	-0.0029 (12)
C32A	0.0482 (18)	0.132 (6)	0.044 (4)	0.040 (2)	0.0079 (16)	-0.020 (4)
C33A	0.051 (3)	0.076 (4)	0.081 (3)	0.016 (3)	0.016 (2)	0.003 (3)
C34A	0.047 (4)	0.133 (7)	0.134 (5)	0.019 (3)	0.023 (4)	-0.014 (5)
C32B	0.0482 (18)	0.132 (6)	0.044 (4)	0.040 (2)	0.0079 (16)	-0.020 (4)
C33B	0.051 (3)	0.076 (4)	0.081 (3)	0.016 (3)	0.016 (2)	0.003 (3)
C34B	0.047 (4)	0.133 (7)	0.134 (5)	0.019 (3)	0.023 (4)	-0.014 (5)

*Geometric parameters (Å, °)*

Cl1—C5	1.726 (2)	C14—H14	0.9300
Cl2—C7	1.739 (2)	C15—H15	0.9300
Cl3—C20	1.730 (3)	C16—C17	1.520 (3)
Cl4A—C22	1.66 (3)	C16—H16	0.9800
Cl4B—C22	1.737 (4)	C17—H17	0.9800
S—C17	1.820 (2)	C18—C23	1.386 (3)
S—C1	1.839 (3)	C18—C19	1.402 (3)
O1—C3	1.379 (3)	C19—C20	1.396 (3)
O1—C1	1.446 (3)	C20—C21	1.381 (3)
O2—C9	1.214 (3)	C21—C22	1.383 (4)
O3—N2	1.207 (4)	C21—H21	0.9300
O4—N2	1.205 (3)	C22—C23	1.370 (4)
O5—C18	1.361 (3)	C23—H23	0.9300
O5—C16	1.430 (3)	C24—C25	1.507 (3)
O6—C24	1.204 (3)	C25—C26	1.373 (3)
O7—N4	1.219 (4)	C25—C30	1.389 (3)
O8—N4	1.214 (3)	C26—C27	1.384 (4)
N1—C9	1.375 (3)	C26—H26	0.9300
N1—C4	1.417 (3)	C27—C28	1.365 (4)
N1—C2	1.468 (3)	C27—H27	0.9300
N2—C13	1.479 (3)	C28—C29	1.369 (4)
N3—C24	1.402 (3)	C29—C30	1.379 (4)
N3—C19	1.422 (3)	C29—H29	0.9300
N3—C17	1.452 (3)	C30—H30	0.9300
N4—C28	1.477 (4)	C31A—C32A	1.495 (7)
C1—C31A	1.519 (3)	C31A—H31A	0.9700
C1—C2	1.568 (3)	C31A—H31B	0.9700
C2—C16	1.522 (3)	C32A—C33A	1.546 (7)
C2—H2	0.9800	C32A—H32A	0.9700
C3—C8	1.381 (3)	C32A—H32B	0.9700
C3—C4	1.387 (3)	C33A—C34A	1.498 (7)
C4—C5	1.395 (3)	C33A—H33A	0.9700
C5—C6	1.388 (3)	C33A—H33B	0.9700
C6—C7	1.378 (4)	C34A—H34A	0.9600
C6—H6	0.9300	C34A—H34B	0.9600

C7—C8	1.376 (4)	C34A—H34C	0.9600
C8—H8	0.9300	C32B—C33B	1.547 (9)
C9—C10	1.495 (3)	C32B—H32C	0.9700
C10—C15	1.375 (3)	C32B—H32D	0.9700
C10—C11	1.393 (3)	C33B—C34B	1.508 (9)
C11—C12	1.371 (4)	C33B—H33C	0.9700
C11—H11	0.9300	C33B—H33D	0.9700
C12—C13	1.363 (4)	C34B—H34D	0.9600
C12—H12	0.9300	C34B—H34E	0.9600
C13—C14	1.371 (4)	C34B—H34F	0.9600
C14—C15	1.381 (3)		
C17—S—C1	93.28 (10)	C16—C17—H17	108.7
C3—O1—C1	113.94 (16)	S—C17—H17	108.7
C18—O5—C16	118.98 (17)	O5—C18—C23	115.5 (2)
C9—N1—C4	126.79 (18)	O5—C18—C19	123.1 (2)
C9—N1—C2	117.17 (18)	C23—C18—C19	121.3 (2)
C4—N1—C2	116.04 (17)	C20—C19—C18	117.2 (2)
O4—N2—O3	124.0 (3)	C20—C19—N3	123.7 (2)
O4—N2—C13	118.1 (3)	C18—C19—N3	118.9 (2)
O3—N2—C13	118.0 (3)	C21—C20—C19	121.8 (2)
C24—N3—C19	121.34 (19)	C21—C20—C13	117.11 (19)
C24—N3—C17	118.40 (19)	C19—C20—C13	121.00 (18)
C19—N3—C17	114.75 (18)	C20—C21—C22	118.8 (2)
O8—N4—O7	124.1 (3)	C20—C21—H21	120.6
O8—N4—C28	118.3 (3)	C22—C21—H21	120.6
O7—N4—C28	117.5 (3)	C23—C22—C21	121.4 (2)
O1—C1—C31A	105.18 (18)	C23—C22—C14A	114.3 (13)
O1—C1—C2	112.17 (17)	C21—C22—C14A	120.4 (11)
C31A—C1—C2	112.8 (2)	C23—C22—C14B	119.6 (2)
O1—C1—S	109.56 (15)	C21—C22—C14B	119.0 (2)
C31A—C1—S	110.25 (18)	C22—C23—C18	119.3 (3)
C2—C1—S	106.90 (15)	C22—C23—H23	120.4
N1—C2—C16	110.95 (18)	C18—C23—H23	120.4
N1—C2—C1	110.10 (16)	O6—C24—N3	122.9 (2)
C16—C2—C1	110.10 (18)	O6—C24—C25	122.5 (2)
N1—C2—H2	108.5	N3—C24—C25	114.5 (2)
C16—C2—H2	108.5	C26—C25—C30	119.5 (2)
C1—C2—H2	108.5	C26—C25—C24	119.6 (2)
O1—C3—C8	120.5 (2)	C30—C25—C24	120.9 (2)
O1—C3—C4	116.83 (19)	C25—C26—C27	120.9 (3)
C8—C3—C4	122.7 (2)	C25—C26—H26	119.6
C3—C4—C5	118.2 (2)	C27—C26—H26	119.6
C3—C4—N1	115.59 (19)	C28—C27—C26	118.2 (3)
C5—C4—N1	125.9 (2)	C28—C27—H27	120.9
C6—C5—C4	119.9 (2)	C26—C27—H27	120.9
C6—C5—C11	119.49 (19)	C27—C28—C29	122.6 (3)
C4—C5—C11	120.52 (18)	C27—C28—N4	119.0 (3)
C7—C6—C5	119.2 (2)	C29—C28—N4	118.3 (3)

C7—C6—H6	120.4	C28—C29—C30	118.6 (3)
C5—C6—H6	120.4	C28—C29—H29	120.7
C8—C7—C6	122.5 (2)	C30—C29—H29	120.7
C8—C7—C12	119.4 (2)	C29—C30—C25	120.2 (3)
C6—C7—C12	118.1 (2)	C29—C30—H30	119.9
C7—C8—C3	117.1 (2)	C25—C30—H30	119.9
C7—C8—H8	121.4	C32A—C31A—C1	116.2 (4)
C3—C8—H8	121.4	C32A—C31A—H31A	108.2
O2—C9—N1	120.2 (2)	C1—C31A—H31A	108.2
O2—C9—C10	120.6 (2)	C32A—C31A—H31B	108.2
N1—C9—C10	119.1 (2)	C1—C31A—H31B	108.2
C15—C10—C11	119.9 (2)	H31A—C31A—H31B	107.4
C15—C10—C9	122.7 (2)	C31A—C32A—C33A	105.4 (5)
C11—C10—C9	117.2 (2)	C31A—C32A—H32A	110.7
C12—C11—C10	120.7 (2)	C33A—C32A—H32A	110.7
C12—C11—H11	119.6	C31A—C32A—H32B	110.7
C10—C11—H11	119.6	C33A—C32A—H32B	110.7
C13—C12—C11	117.8 (3)	H32A—C32A—H32B	108.8
C13—C12—H12	121.1	C34A—C33A—C32A	106.9 (5)
C11—C12—H12	121.1	C34A—C33A—H33A	110.3
C12—C13—C14	123.3 (2)	C32A—C33A—H33A	110.4
C12—C13—N2	117.7 (3)	C34A—C33A—H33B	110.4
C14—C13—N2	119.0 (3)	C32A—C33A—H33B	110.4
C13—C14—C15	118.4 (2)	H33A—C33A—H33B	108.6
C13—C14—H14	120.8	C33B—C32B—H32C	105.8
C15—C14—H14	120.8	C33B—C32B—H32D	105.8
C10—C15—C14	119.8 (2)	H32C—C32B—H32D	106.2
C10—C15—H15	120.1	C34B—C33B—C32B	110.4 (11)
C14—C15—H15	120.1	C34B—C33B—H33C	109.6
O5—C16—C17	111.36 (18)	C32B—C33B—H33C	109.6
O5—C16—C2	105.82 (17)	C34B—C33B—H33D	109.6
C17—C16—C2	107.66 (17)	C32B—C33B—H33D	109.6
O5—C16—H16	110.6	H33C—C33B—H33D	108.1
C17—C16—H16	110.6	C33B—C34B—H34D	109.5
C2—C16—H16	110.6	C33B—C34B—H34E	109.5
N3—C17—C16	111.85 (17)	H34D—C34B—H34E	109.5
N3—C17—S	113.76 (16)	C33B—C34B—H34F	109.5
C16—C17—S	105.00 (15)	H34D—C34B—H34F	109.5
N3—C17—H17	108.7	H34E—C34B—H34F	109.5
C3—O1—C1—C31A	173.0 (2)	N1—C2—C16—C17	-88.4 (2)
C3—O1—C1—C2	50.0 (2)	C1—C2—C16—C17	33.7 (2)
C3—O1—C1—S	-68.5 (2)	C24—N3—C17—C16	-107.8 (2)
C17—S—C1—O1	107.13 (15)	C19—N3—C17—C16	46.4 (3)
C17—S—C1—C31A	-137.58 (17)	C24—N3—C17—S	133.40 (18)
C17—S—C1—C2	-14.64 (15)	C19—N3—C17—S	-72.4 (2)
C9—N1—C2—C16	-97.4 (2)	O5—C16—C17—N3	-51.8 (2)
C4—N1—C2—C16	81.7 (2)	C2—C16—C17—N3	-167.39 (18)
C9—N1—C2—C1	140.5 (2)	O5—C16—C17—S	72.02 (19)

C4—N1—C2—C1	-40.5 (2)	C2—C16—C17—S	-43.55 (19)
O1—C1—C2—N1	-6.0 (3)	C1—S—C17—N3	156.25 (16)
C31A—C1—C2—N1	-124.5 (2)	C1—S—C17—C16	33.63 (15)
S—C1—C2—N1	114.15 (17)	C16—O5—C18—C23	175.8 (2)
O1—C1—C2—C16	-128.59 (19)	C16—O5—C18—C19	-6.2 (3)
C31A—C1—C2—C16	112.9 (2)	O5—C18—C19—C20	-176.4 (2)
S—C1—C2—C16	-8.5 (2)	C23—C18—C19—C20	1.6 (4)
C1—O1—C3—C8	131.2 (2)	O5—C18—C19—N3	-0.7 (4)
C1—O1—C3—C4	-48.9 (3)	C23—C18—C19—N3	177.2 (2)
O1—C3—C4—C5	174.5 (2)	C24—N3—C19—C20	-51.9 (3)
C8—C3—C4—C5	-5.6 (3)	C17—N3—C19—C20	154.7 (2)
O1—C3—C4—N1	0.1 (3)	C24—N3—C19—C18	132.7 (2)
C8—C3—C4—N1	-180.0 (2)	C17—N3—C19—C18	-20.7 (3)
C9—N1—C4—C3	-134.9 (2)	C18—C19—C20—C21	-4.1 (4)
C2—N1—C4—C3	46.1 (3)	N3—C19—C20—C21	-179.5 (2)
C9—N1—C4—C5	51.2 (3)	C18—C19—C20—C13	173.25 (18)
C2—N1—C4—C5	-127.8 (2)	N3—C19—C20—C13	-2.2 (3)
C3—C4—C5—C6	6.7 (3)	C19—C20—C21—C22	3.8 (4)
N1—C4—C5—C6	-179.6 (2)	C13—C20—C21—C22	-173.7 (2)
C3—C4—C5—C11	-169.92 (17)	C20—C21—C22—C23	-0.8 (4)
N1—C4—C5—C11	3.8 (3)	C20—C21—C22—C14A	156 (3)
C4—C5—C6—C7	-3.3 (4)	C20—C21—C22—C14B	177.8 (3)
C11—C5—C6—C7	173.38 (19)	C21—C22—C23—C18	-1.6 (5)
C5—C6—C7—C8	-1.6 (4)	C14A—C22—C23—C18	-160 (3)
C5—C6—C7—C12	-178.06 (19)	C14B—C22—C23—C18	179.7 (3)
C6—C7—C8—C3	2.8 (4)	O5—C18—C23—C22	179.3 (3)
C12—C7—C8—C3	179.19 (18)	C19—C18—C23—C22	1.2 (4)
O1—C3—C8—C7	-179.2 (2)	C19—N3—C24—O6	-9.3 (3)
C4—C3—C8—C7	0.9 (4)	C17—N3—C24—O6	143.2 (2)
C4—N1—C9—O2	-165.7 (2)	C19—N3—C24—C25	168.6 (2)
C2—N1—C9—O2	13.3 (3)	C17—N3—C24—C25	-39.0 (3)
C4—N1—C9—C10	17.7 (3)	O6—C24—C25—C26	-48.0 (4)
C2—N1—C9—C10	-163.4 (2)	N3—C24—C25—C26	134.1 (3)
O2—C9—C10—C15	-130.6 (3)	O6—C24—C25—C30	129.3 (3)
N1—C9—C10—C15	46.1 (4)	N3—C24—C25—C30	-48.5 (3)
O2—C9—C10—C11	43.3 (4)	C30—C25—C26—C27	-0.3 (4)
N1—C9—C10—C11	-140.1 (3)	C24—C25—C26—C27	177.1 (3)
C15—C10—C11—C12	-0.9 (4)	C25—C26—C27—C28	-1.0 (5)
C9—C10—C11—C12	-175.0 (3)	C26—C27—C28—C29	0.6 (5)
C10—C11—C12—C13	1.4 (5)	C26—C27—C28—N4	179.2 (3)
C11—C12—C13—C14	-1.1 (5)	O8—N4—C28—C27	167.9 (3)
C11—C12—C13—N2	-179.8 (3)	O7—N4—C28—C27	-13.4 (5)
O4—N2—C13—C12	173.8 (3)	O8—N4—C28—C29	-13.5 (4)
O3—N2—C13—C12	-7.3 (5)	O7—N4—C28—C29	165.2 (3)
O4—N2—C13—C14	-4.9 (5)	C27—C28—C29—C30	1.1 (5)
O3—N2—C13—C14	174.0 (3)	N4—C28—C29—C30	-177.5 (3)
C12—C13—C14—C15	0.3 (5)	C28—C29—C30—C25	-2.4 (5)
N2—C13—C14—C15	179.0 (3)	C26—C25—C30—C29	2.0 (4)
C11—C10—C15—C14	0.1 (4)	C24—C25—C30—C29	-175.3 (3)

C9—C10—C15—C14	173.8 (2)	O1—C1—C31A—C32A	-44.4 (6)
C13—C14—C15—C10	0.2 (4)	C2—C1—C31A—C32A	78.1 (6)
C18—O5—C16—C17	31.9 (3)	S—C1—C31A—C32A	-162.5 (5)
C18—O5—C16—C2	148.64 (19)	C1—C31A—C32A—C33A	174.9 (4)
N1—C2—C16—O5	152.39 (17)	C31A—C32A—C33A—C34A	-178.8 (7)
C1—C2—C16—O5	-85.5 (2)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C2—H2 $\cdots$ O2 <sup>i</sup>	0.98	2.55	3.491 (3)	162
C14—H14 $\cdots$ O3 <sup>ii</sup>	0.93	2.41	3.298 (4)	160

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $-x+1, y+1/2, -z+3/2$ .